

## Incorporation of DNA and Protein Precursors into Macromolecules by Bacteria at $-15^{\circ}\text{C}$

Brent C. Christner\*

Department of Microbiology, The Ohio State University, Columbus, Ohio 43210-1292

Received 24 June 2002/Accepted 30 August 2002

**DNA and protein precursors were incorporated into trichloroacetic acid-precipitated material by bacterial cell suspensions during incubation for 50 to 100 days at  $-15^{\circ}\text{C}$ . Incorporation did not occur at  $-70^{\circ}\text{C}$  and was inhibited by antibiotics. The results demonstrate that bacteria can perform macromolecular synthesis under conditions that mimic entrapment in glacial ice.**

Bacteria can remain viable for hundreds of thousands of years when trapped in glacial ice (1, 2, 8 [http://www.ohiolink.edu/etd/send-pdf.cgi?osu1015965965], 9). In the absence of repair, macromolecular damage must accumulate through amino acid racemization, DNA depurination, and exposure to natural ionizing radiation (12, 13). However, it is known that temperature and the presence of water strongly influence macromolecular decay and that protein and nucleic acid decay rates are drastically reduced within materials with low water activity, such as amber (4) and perhaps ice as well. It also seems possible that such entrapped microbes might carry out slow rates of metabolism to repair the incurred macromolecular damage. Thin veins of liquid exist between ice crystals that could provide a habitat for microorganisms within apparently solid ice (15), and studies of permafrost (17) and South Pole snow (6) have detected low levels of metabolic activity at sub-zero temperatures. Therefore, to explore the concept that microorganisms trapped in glacial ice might also be metabolically active, reconstruction experiments were undertaken to determine if macromolecular synthesis occurs at  $-15^{\circ}\text{C}$  when bacteria isolated from glacial ice core samples were refrozen.

**Bacterial strains and culture conditions.** A  $\gamma$ -proteobacterial *Psychrobacter* species (8) (isolate Trans1; 16S ribosomal DNA [rDNA] GenBank accession no. AF479327) and *Escherichia coli* (Ohio State reference no. 422) were grown in Luria-Bertani medium (18), and an actinobacterial *Arthrobacter* species (7) (isolate G200-C1; 16S rDNA GenBank accession no. AF479341) was cultured in R2 medium (16). Cultures (25 ml) were incubated aerobically with shaking (200 rpm) at  $22^{\circ}\text{C}$  in 125-ml Erlenmeyer flasks to the late exponential growth phase and were diluted to an initial  $A_{600}$  of 0.2 and allowed to grow to an  $A_{600}$  of 0.6. The cells present were then harvested by centrifugation at  $17,000 \times g$  for 5 min, washed twice with distilled water, and resuspended at an  $A_{600}$  of 0.2 in distilled water (equivalent to  $1.3 \times 10^8$  and  $2 \times 10^8$  CFU  $\text{ml}^{-1}$  for Trans1 and G200-C1, respectively). Aliquots (500  $\mu\text{l}$ ) of these cell suspensions were placed in 1.5-ml Eppendorf tubes and chilled to  $4^{\circ}\text{C}$ .

**Precursor incorporation assays.** Cell suspensions were maintained on ice, and 100  $\mu\text{l}$  of a chilled working solution (10  $\mu\text{Ci ml}^{-1}$ ; diluted 1:100 from stock) of either [ $^3\text{H}$ ]thymidine (ICN Biomedicals, catalog no. 24060; 60 to 90 Ci/mmol in sterile water) or [ $^3\text{H}$ ]leucine (ICN Biomedicals, catalog no. 20036E; 40 to 60 Ci/mmol in a sterile 2:98 ethanol-water mixture) was added to each sample, and the mixture was rapidly frozen by incubation at  $-70^{\circ}\text{C}$ . After 1 h at  $-70^{\circ}\text{C}$ , tubes were transferred to a  $-15^{\circ}\text{C}$  freezer, except for control cell suspensions that were maintained at  $-70^{\circ}\text{C}$ . At designated experimental time points, 100  $\mu\text{l}$  of 50% trichloroacetic acid (TCA) was added to a frozen mixture, which was then allowed to melt at  $4^{\circ}\text{C}$ . After 30 min at  $4^{\circ}\text{C}$ , the TCA-insoluble macromolecules were sedimented by centrifugation at  $18,000 \times g$  for 15 min. The pellet was washed with 500  $\mu\text{l}$  of 5% TCA, recentrifuged for 10 min at  $18,000 \times g$ , washed with 500  $\mu\text{l}$  of 70% ethanol, and suspended in 1 ml of Ecoscint H scintillation fluid (Life Sciences, Inc., catalog no. LS-275). The Eppendorf tube was placed into a scintillation vial, and the radioactivity present was quantitated by liquid scintillation counting (Beckman model LS-7500 scintillation counter).

**Incorporation into TCA-precipitable material at  $-15^{\circ}\text{C}$ .** [ $^3\text{H}$ ]thymidine and [ $^3\text{H}$ ]leucine were incorporated into TCA-precipitated material by cells frozen in distilled water during incubation at  $-15^{\circ}\text{C}$  (Fig. 1). The majority of the incorporation occurred within the first 50 days after freezing. Incorporation did not occur in identical samples incubated at  $-70^{\circ}\text{C}$ , a temperature below that predicted for liquid water to exist in ice (14), or by controls pretreated with 5% TCA before freezing. The radioactive counts observed in TCA-treated controls were similar in samples incubated at  $-15$  and  $22^{\circ}\text{C}$  (1,000 to 5,000 dpm), indicating that  $\sim 0.2\%$  of the radiolabel persisted following  $\sim 1:1,000$  dilution of the mixture during precipitation and washing. Since there is not a significant difference between the radioactive counts obtained in samples prefixed in TCA and those incubated at  $-70^{\circ}\text{C}$  in distilled water, this suggests that the background in these experiments results from persistence of residual radiolabel and is not due to acid conditions effecting chemical binding.

No precursors were incorporated after  $>5$  months at  $-15^{\circ}\text{C}$  by cells frozen in their respective growth media (data not shown). The presence of substrate in these undefined media may have diluted the radiolabel and prevented detection of a

\* Present address: Land Resources & Environmental Sciences, Montana State University, 334 Leon Johnson Hall, Bozeman, MT 59717. Phone: (406) 994-2733. Fax: (406) 994-5863. E-mail: bchristner@montana.edu.




FIG. 1. Incorporation of [<sup>3</sup>H]thymidine and [<sup>3</sup>H]leucine into TCA-precipitable material over 172 days by frozen suspensions of *Psychrobacter* sp. strain Trans1 and *Arthrobacter* sp. strain G200-C1 at -15 and -70°C. Duplicate samples were measured at each time point. The y-axis error bars denote standard deviations from the mean.

low level of incorporation over this period of time. Another explanation is that nutrient-fed cultures were not metabolically stressed prior to freezing, as would occur when cells in logarithmic growth were washed and reconstituted in distilled water. During metabolic arrest, the imbalance between decreased anabolism and continuing catabolism causes a burst of free radical production, resulting in damage to DNA and other cellular components (3). It is possible that incorporation by cells in ice made from distilled water represents activity directed toward repair of damage incurred prior to freezing.

In experiments in which the viable count was also determined, the recovery of G200-C1 decreased fourfold during the freezing process and then an additional fourfold during the first 12 days of incubation at -15°C (Fig. 2). Due to a technical

error, a day 0 time point is not available for the Trans1 isolate, but the viable count was 12-fold lower after 2 days at -15°C than before freezing. In both cases, the numbers of CFU per milliliter recovered stabilized by 150 days postfreezing. Interestingly, 70 to 80% of the thymidine incorporation occurred during the first 12 days of incubation, and this coincided with the period of time when the numbers of CFU per milliliter became stable, after the initial 17-fold decrease that occurred during freezing and initial storage for 12 days at -15°C. The presence of ciprofloxacin (ICN Biomedicals, catalog no. 199020) reduced thymidine incorporation by 30 to 40%, and the presence of chloramphenicol (Sigma, catalog no. 100K9113) reduced leucine incorporation 50 to 60% (Table 1). Curiously, incubations in the presence of nalidixic acid (Sigma,




FIG. 2. Incorporation of  $[^3\text{H}]$ thymidine and numbers of CFU per milliliter for *Psychrobacter* sp. strain Trans1 and *Arthrobacter* sp. strain G200-C1 over 274 days of incubation at  $-15^{\circ}\text{C}$ . Before freezing, the cell suspensions of Trans1 and G200-C1 contained  $1.3 \times 10^8$  CFU  $\text{ml}^{-1}$  and  $2 \times 10^8$  CFU  $\text{ml}^{-1}$ , respectively. The 2-day number of CFU per milliliter represents the first data point for Trans1. Each point shown was the result of triplicate measurements. The y-axis error bars denote standard deviations from the mean.

catalog no. N-3143), resulted in a twofold increase in thymidine incorporation in both species, even though growth of G200-C1 is inhibited in the presence of  $15 \mu\text{g}$  of this antibiotic per ml. Precursor incorporation was similarly affected in aqueous cell suspensions incubated at  $22^{\circ}\text{C}$  for 20 h in the presence of chloramphenicol and nalidixic acid (data not shown). Although the amount of incorporation under liquid conditions after 20 h at  $22^{\circ}\text{C}$  was substantially larger (10 to  $100\times$ ) than after 23 days at  $-15^{\circ}\text{C}$ , leucine incorporation was 25 to 85% lower in the presence of chloramphenicol, and thymidine incorporation increased twofold in the presence of nalidixic acid.

**Biochemical fractionation.** The washed, TCA-precipitated material from cells incubated at  $-15^{\circ}\text{C}$  for 280 days with  $[^3\text{H}]$ thymidine or  $[^3\text{H}]$ leucine was fractionated according to the method of Kelley (11). This material was suspended in  $30 \mu\text{l}$  of a 1:1 ethanol-ether mixture, incubated at  $37^{\circ}\text{C}$  for 30 min,

and then centrifuged for 10 min at  $18,000 \times g$ . The resulting supernatant was collected, and the pellet was resuspended in  $30 \mu\text{l}$  of 5% TCA and incubated at  $95^{\circ}\text{C}$  for 30 min. After centrifugation at  $18,000 \times g$  for 10 min, the supernatant was collected, and the radioactivity in all fractions was quantitated by liquid scintillation counting.

In both cases,  $\sim 50\%$  of the  $[^3\text{H}]$ thymidine was incorporated into the hot TCA extract containing nucleic acids, and  $\sim 50\%$  was incorporated into the insoluble residue that would contain proteins (Table 2). Although less  $[^3\text{H}]$ leucine was incorporated, the majority of this precursor was incorporated into the protein fraction. Given that 7,400 to 8,400 dpm of thymidine and  $\sim 2,100$  dpm of leucine were incorporated per sample into nucleic acid and protein, respectively, it can be calculated from the specific activities of the precursors (60 to 90 Ci of  $[^3\text{H}]$ thymidine and 40 to 60 Ci of  $[^3\text{H}]$ leucine per mmol) and from the

TABLE 1. Incorporation of [<sup>3</sup>H]thymidine and [<sup>3</sup>H]leucine by frozen cell suspensions during 23 days of incubation at -15°C<sup>a</sup>

Experimental parameter	Precursor incorporation into TCA-precipitated material (dpm)
<i>Psychrobacter</i> sp. strain Trans1	
[ <sup>3</sup> H]thymidine	
No inhibitor.....	8,500 ± 1,200
5% TCA.....	4,400 ± 190
+ Ciprofloxacin.....	4,900 ± 640
+ Nalidixic acid.....	10,000 ± 1,600
[ <sup>3</sup> H]leucine	
No inhibitor.....	5,000 ± 330
5% TCA.....	3,500 ± 330
+ Chloramphenicol.....	2,400 ± 330
<i>Arthrobacter</i> sp. strain G200-A1	
[ <sup>3</sup> H]thymidine	
No inhibitor.....	11,000 ± 300
5% TCA.....	3,000 ± 290
+ Ciprofloxacin.....	7,900 ± 170
+ Nalidixic acid.....	22,000 ± 400
[ <sup>3</sup> H]leucine	
No inhibitor.....	8,800 ± 1,300
5% TCA.....	2,600 ± 170
+ Chloramphenicol.....	3,100 ± 35

<sup>a</sup> As indicated, 5% TCA (final concentration) or 15 µg of the antibiotic listed per ml was added to the cell suspension before the addition of radioactive precursor. The values are means ± standard deviations (*n* = 2).

assumption that all cells present ( $1.1 \times 10^7$  CFU ml<sup>-1</sup>) (Fig. 2) participated in the DNA and protein syntheses that 4,100 to 6,900 molecules of thymidine and ~2,000 molecules of leucine were incorporated per cell after 280 days of incubation. Based on 3-Mbp genomes and an average protein size of 36 kDa, this incorporation corresponds to replication of <1% of the genome and synthesis of ~100 protein molecules. This incorporation is therefore insufficient for reproduction.

**Implications for metabolism in glacial ice.** The results are consistent with macromolecular synthesis at -15°C by isolates recovered from polar and nonpolar glacial ice cores under conditions comparable to those within glacial ice. The 16S rDNA sequences of these psychrotrophic isolates are >99% identical to those of species recovered from brine channels in sea ice (5, 10), and therefore it was hypothesized that a biological activity at below-freezing temperatures might be exclusive to closely related strains. However, [<sup>3</sup>H]thymidine and

TABLE 2. Fractionation of TCA-insoluble material from cell suspensions incubated at -15°C in the presence of [<sup>3</sup>H]thymidine and [<sup>3</sup>H]leucine for 280 days<sup>a</sup>

Macromolecular fraction ( <i>n</i> = 2)	<i>Psychrobacter</i> sp. strain Trans1		<i>Arthrobacter</i> sp. strain G200-A1	
	[ <sup>3</sup> H]thymidine incorporation (dpm)	[ <sup>3</sup> H]leucine incorporation (dpm)	[ <sup>3</sup> H]thymidine incorporation (dpm)	[ <sup>3</sup> H]leucine incorporation (dpm)
Total TCA-insoluble material	18,000 ± 650	3,400 ± 320	16,000 ± 350	2,600 ± 520
Ethanol-ether soluble (lipid)	930 ± 82	1,200 ± 280	1,100 ± 400	680 ± 180
Hot 5% TCA soluble (nucleic acid)	8,400 ± 860	1,800 ± 160	7,400 ± 26	1,600 ± 64
Residual (protein)	8,200 ± 1,100	2,100 ± 54	11,000 ± 1,800	2,100 ± 340

<sup>a</sup> Values are means ± standard deviations (*n* = 2).

[<sup>3</sup>H]leucine were also incorporated into TCA-precipitable material during 102 days of incubation at -15°C by a laboratory strain of *E. coli*, indicating that this ability is not limited to species specialized for survival in ice. For cells in 5% TCA, 3,800 ± 1,100 dpm (mean ± standard deviation) of [<sup>3</sup>H]thymidine and 4,100 ± 16 dpm of [<sup>3</sup>H]leucine were incorporated (*n* = 2). For cells in distilled water, 15,000 ± 470 dpm of [<sup>3</sup>H]thymidine and 17,000 ± 9,900 dpm of [<sup>3</sup>H]leucine were incorporated (*n* = 2).

The data reported here add to the growing body of evidence for metabolism under frozen conditions (6, 17). They support the argument that bacteria trapped in glacial ice might repair macromolecular damage that occurs while immured for extended periods. Possibly, cells remain metabolically active in solute-enriched water films on the surface of entrapped particulates and air bubbles or in liquid veins between adjacent ice crystals (15).

I am indebted to John Reeve for experimental guidance and valuable editorial comments and thank Wade Jeffrey for providing procedural advice.

This research was supported at Ohio State University by National Science Foundation (NSF) grant OPP-9714206, awarded through the Life in Extreme Environments Initiative. I am presently supported at Montana State University by NSF grant OPP-0085400.

## REFERENCES

1. **Abyzov, S. S.** 1993. Microorganisms in the Antarctic ice, p. 265–295. In E. I. Friedmann (ed.), *Antarctic microbiology*. Wiley-Liss, Inc., New York, N.Y.
2. **Abyzov, S. S., I. N. Mitskevich, and M. N. Poglazova.** 1998. Microflora of the deep glacier horizons of central Antarctica. *Microbiology (Moscow)* **67**:66–73.
3. **Aldsworth, T. G., R. L. Sharman, and C. E. R. Dodd.** 1999. Bacterial suicide through stress. *Cell. Mol. Life Sci.* **56**:378–383.
4. **Bada, J. L., X. S. Wang, H. N. Poinar, S. Paabo, and G. O. Poinar.** 1994. Amino acid racemization in amber-entombed insects: implications for DNA preservation. *Geochim. Cosmochim. Acta* **58**:3131–3135.
5. **Bowman, J. P., S. A. McCammon, M. V. Brown, D. S. Nichols, and T. A. McMeekin.** 1997. Diversity and association of psychrophilic bacteria in Antarctic sea ice. *Appl. Environ. Microbiol.* **63**:3068–3078.
6. **Carpenter, E. J., S. Lin, and D. G. Capone.** 2000. Bacterial activity in South Pole snow. *Appl. Environ. Microbiol.* **66**:4514–4517.
7. **Christner, B. C., E. Mosley-Thompson, L. G. Thompson, V. Zagorodnov, K. Sandman, and J. N. Reeve.** 2000. Recovery and identification of viable bacteria immured in glacial ice. *Icarus* **144**:479–485.
8. **Christner, B. C.** 2002. Detection, recovery, isolation, and characterization of bacteria in glacial ice and Lake Vostok accretion ice. Ph.D. thesis. The Ohio State University, Columbus.
9. **Christner, B. C., E. Mosley-Thompson, L. G. Thompson, and J. N. Reeve.** Bacterial recovery from ancient glacial ice. *Environ. Microbiol.*, in press.
10. **Junge, K., J. J. Gosink, H. G. Hoppe, and J. T. Staley.** 1998. *Arthrobacter*, *Brachybacterium* and *Planococcus* isolates identified from Antarctic sea ice brine. Description of *Planococcus mceekini*, sp. nov. *Syst. Appl. Microbiol.* **21**:306–314.
11. **Kelley, D. P.** 1967. The incorporation of acetate by the chemoautotroph *Thiobacillus neapolitanus* strain C. *Arch. Mikrobiol.* **58**:99–116.
12. **Kennedy, M. J., S. L. Reader, and L. M. Swierczynski.** 1994. Preservation records of micro-organisms: evidence of the tenacity of life. *Microbiology* **140**:2513–2529.
13. **McKay, C. P.** 2001. The deep biosphere: lessons for planetary exploration, p. 315–327. In J. K. Fredrickson and M. Fletcher (ed.), *Subsurface microbiology and biogeochemistry*. Wiley-Liss, Inc., New York, N.Y.
14. **Ostromov, V. E., and C. Siegert.** 1996. Exobiological aspects of mass transfer in microzones of permafrost deposits. *Adv. Space Res.* **18**:79–86.
15. **Price, B. P.** 2000. A habitat for psychrophiles in deep Antarctic ice. *Proc. Natl. Acad. Sci. USA* **97**:1247–1251.
16. **Reasoner, D. J., and E. E. Geldreich.** 1985. A new medium for the enumeration and subculture of bacteria from potable water. *Appl. Environ. Microbiol.* **49**:1–7.
17. **Rivkina, E. M., E. I. Friedmann, C. P. McKay, and D. A. Gilichinsky.** 2000. Metabolic activity of permafrost bacteria below the freezing point. *Appl. Environ. Microbiol.* **66**:3230–3233.
18. **Sambrook, J., E. F. Fritsch, and T. Maniatis.** 1989. *Molecular cloning: a laboratory manual*, 2nd ed. Cold Spring Harbor Press, Cold Spring Harbor, New York.